



Original Article

Bleeding in Pancreatitis and its management

Dr. Hiranmoy Das¹, Dr. Mubashshir Ahmed²

¹ Assistant Professor, Department of Surgery, Gauhati Medical College, Guwahati, Assam

² Assistant Professor, Department of Surgery, Nalbari Medical College, Nalbari, Assam

OPEN ACCESS

Corresponding Author:

Dr. Hiranmoy Das

Assistant Professor, Department of
Surgery, Gauhati Medical College,
Guwahati, Assam.

Received: 07-04-2026

Accepted: 02-05-2026

Available online: 25-05-2026

Copyright © International Journal of
Medical and Pharmaceutical Research

ABSTRACT

BACKGROUND: Pancreatitis is associated with arterial complications in 4-10% of patients. If untreated, mortality approaches 90%. Timely intervention can reduce the mortality to 15%. We present a single institution experience of open or selective embolization as first line management of bleeding pseudoaneurysms in pancreatitis. **AIM:** This Retrospective, hospital based observational study conducted at a tertiary care centre in Assam aims to evaluate the incidence of mortality and rebleeding in patients of severe pancreatitis with bleeding, in cases undergoing open surgery and angioembolization.

METHODS & MATERIALS: 34 patients with pancreatitis and bleeding were identified in a retrospective study from January 2019 to December 2024 from surgical records at Gauhati Medical College & Hospital, Dept. of Surgery. True visceral artery aneurysm and pseudoaneurysm arising as a result of post operative pancreatic & biliary leak were excluded from the study.

RESULTS: Patients were divided into acute & chronic pancreatitis group. All patients had a contrast CT abdomen scan & 12 patients underwent celiac axis angiography. The bleeding affected the splenic artery in 12 patients, Gastroduodenal-8, Superior Pancreatic artery-2, left gastric -3, pancreaticoduodenal-1 and rest cavity bleeding (source not identified). In the acute group, 14 cases underwent surgery while 6 cases had angioembolization. Mortality rate was 35% in surgery group & 17% in angioembolization group. While in the chronic pancreatitis group, 3 cases underwent surgery and 11 cases had angioembolization. Mortality in Angioembolization group was 9%.

CONCLUSIONS: In pancreatitis, rebleeding & mortality rates are less in angio embolization of visceral vessels than open surgery.

Keywords: Pseudoaneurysms, Pancreatitis, Vascular complications, surgery, Embolization.

INTRODUCTION

Pancreatitis is associated with arterial complications in 4-10% of patients, which if untreated mortality approaches 90%. Timely intervention can reduce mortality to 15%. Pathophysiologically; progressive inflammation within pancreas and surrounding tissues promotes vessel erosion and formation of pseudoaneurysms by extravasation of proteolytic enzymes, formation of pseudocysts, extensive necrosis and abscesses. They may rupture & bleed either into pseudocyst, peritoneal cavity, retroperitoneum or erosions into stomach, pancreatic duct (hemorrhage pancreatitis) or bile ducts (hemobilia). We present a single institution experience of surgery and angioembolization as first line of management in management of bleeding in pancreatitis.

METHODS

Records of 325 patients with pancreatitis were retrospectively reviewed from January 2019 to December 2024 from department of surgery, GMCH, Guwahati, Assam. Of these, 268 were acute cases and 57 were chronic cases with an age range of 9-84yrs (Mean= 46.5yrs) & 9-78 yrs (Mean=43.5Yrs) respectively. Diagnosis was based on USG, CECT and Angiography. Surgery was the primary mode of treatment prior to September 2020, while surgery or angioembolization was performed thereafter depending on patient condition & hospital availability of interventional radiology expertise.

Median follow-up time of the patients was 6 months. True Visceral artery aneurysm and Pseudoaneurysms due to postoperative cause or bile leak were excluded from the study.

CHARACTERISTICS OF ACUTE GROUP:

20 patients of the acute group with a mean age of 46 years (range= 9-84) had bleeding. 14 of them underwent surgery while 6 patients had angioembolization. Of the surgical group, 7 patients had diffuse bleeding into the necrosectomy cavity (cavity bleed), 4 from splenic artery, 2 from gastroduodenal artery (GDA) & 1 from superior mesenteric artery (SMA). In embolization group, 2 bled from SA, 3 from GDA and 1 from SMA.

CHARACTERISTICS OF THE chronic group:

14 patients of the chronic group with a mean age of 41 years (range= 25-68) had visceral bleeding. Of them, 3 had surgery (2 from SA & 1 from GDA) while 11 patients had angio embolization (4 from SA, 3 from GDA, 3 LGA & 1 from inferior pancreatico duodenal)

Most common symptoms were:

Abdominal Pain (Acute-17, Chronic-17)

Vomiting (acute-15, chronic -19)

Shock (acute-5, chronic-5)

Upper GI bleeding (acute-5, chronic- 4); Malena (acute-15, chronic-12)

MANAGEMENT PROTOCOL:

For surgical exploration, an uniform approach was followed. A midline supra umbilical incision was given & lesser sac entered by dividing the gastrocolic ligament. Necrotic Materials were removed bluntly aided with saline washes. Visible bleeding Vessel was controlled with ligation or in combination with distal pancreatectomy. Uncontrolled bleeding was treated with packing and reassessment after 48hrs.




Fig a. Pseudoaneurysm at splenic artery **b.** Reconstructed angiographic image of the same patient **c.** Digital subtraction angiography of the same patient showing contrast outpouching



Fig: Intraoperative surgical Necrosectomy

For angio embolization, access was taken through a femoral puncture by interventional radiologist and SMA, Coeliac axis and portal veins were imaged on angiography. A micro catheter was used to selectively catheterize the pseudoaneurysm. Super selective embolization was performed with stainless steel coils (0.18) alone or in combination with polyvinyl alcohol(PVA). Embolization was regarded as successful when cessation of bleeding was demonstrated radiologically.

RESULT

In the acute group, 14 cases underwent surgery while 6 had embolization. Necrosectomy and suture ligation of bleeding vessels were done in open surgery. In the surgery group, 7 patients had cavitory bleed (no vessel identified), 4 from SMA, 2 from GDA and 1 from SMA. Four of the cavitory bleeding had rebleed, which were repacked. Total of 7 patients had rebleed(4-cavitory, 2-SA & 1 SMA), and all 7 expired. Mortality rate was 35%. While in the angio embolization group, there were total of 6 patients. Success rate in angioembolization was 83% and mortality was 17%.

In the chronic group, 3 cases underwent surgery and 11 cases had angio-embolization. 2 of the patients that underwent surgery had rebleeding from SA and 1 from GDA. 2 of them had rebleeding and expired. While in the angioembolization group, embolization with coil was done in 10 patients (4-SA, 3-GDA, 3-LGA & 1 Inferior pancreatico duodenal artery). In 1 patient, both coil & PVA was used. In this group, bleeding stopped in 10 patients while 1 patient had rebleeding & expired. Mortality in angio embolization was 9 % and success rate was 91%.

DISCUSSION

Most common clinical presentation in our series included abdominal pain, vomiting, shock & GIT bleeding. Due to low sensitivity of USG abdomen in diagnosing pancreatic bleeding, complications is often ignored even with contrast enhanced USG. CECT abdomen is considered gold standard to evaluate the degree of severity in pancreatitis.

Distribution of visceral artery bleeding has been fairly consistent in literature (SA-40%, GDA-30%, PDA-20%, Gastric artery- 5% & hepatic artery- 2%). Our series also found a similar result with bleeding from SA-39.2%, GDA-26.5%, LGA-5.8%, SMA-5.8% & PDA- 2.9%. It was also seen that source of bleeding are more likely to be identified in angiography (100%) than in surgery (79.4%). The reported success of embolization is 79-100% and the reported mortality rate after embolization is 12-33%, in patients with acute and chronic pancreatic bleeding. Laparotomy for therapy of bleeding should be considered only in haemodynamically unstable patients or unsuccessful coiling. Surgical packing is the procedure of choice in diffuse venous intraabdominal bleeding. In most cases, it seems that angio embolization is sufficient to control bleeding complications. Re-bleeding rate is more in surgical group (35%). Insufficient ligation of a vessel due to surrounding tissue infection, oedema or insufficient control may cause rebleeding.

Mortality seems to be higher in patients of pancreatitis complicated by intra abdominal hemorrhage. In some studies, there is three fold higher probability of a fatal outcome in patients with severe pancreatitis associated with intra abdominal hemorrhage than those with severe pancreatitis without hemorrhagic complications. However, others suggest that hemorrhagic complication per se have little influence on mortality. Bleeding complications seem to reflect the severity of disease than being the direct cause of mortality.

CONCLUSION

Angiography is valuable in localizing & controlling visceral artery bleeding. Rebleeding & mortality rates are less in angio embolization of visceral vessels, than surgery.

REFERENCES

1. Mendelson RM, Anderson J, Marshall M, Ramsay D. Vascular complications of pancreatitis. *ANZ J. Surg.* 2005; 75: 1073–9. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
2. Balachandra S, Siriwardena AK. Systematic appraisal of the management of the major vascular complications of pancreatitis. *Am. J. Surg.* 2005; 190: 489–95. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
3. Bergert H, Hinterseher I, Kersting S, Leonhardt J, Bloomenthal A, Saeger HD. Management and outcome of hemorrhage due to arterial pseudoaneurysms in pancreatitis. *Surgery.* 2005; 137: 323–8. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
4. Morteale KJ, Mergo PJ, Taylor HM *et al* Peripancreatic vascular abnormalities complicating acute pancreatitis: contrast-enhanced helical CT findings. *Eur. J. Radiol.* 2004; 52: 67–72. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
5. Boudghene F, L'Hermine C, Bigot JM. Arterial complications of pancreatitis: diagnostic and therapeutic aspects in 104 cases. *J. Vasc. Interv. Radiol.* 1993; 4: 551–8. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
6. Banks PA, Bollen TL, Dervenis C *et al* Classification of acute pancreatitis—2012: revision of the Atlanta classification and definitions by international consensus. *Gut.* 2013; 62: 102–11. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
7. Singer MV, Gyr K, Sarles H. Revised classification of pancreatitis. Report of the Second International Symposium on the Classification of Pancreatitis in Marseille, France, March 28–30, 1984. *Gastroenterology.* 1985; 89: 683–5. [\[PubMed\]](#) [\[Google Scholar\]](#)
8. Cardella JF, Kundu S, Miller DL, Millward SF, Sacks D, Society of Interventional Radiology . Society of Interventional Radiology clinical practice guidelines. *J. Vasc. Interv. Radiol.* 2009; 20(7 Suppl): S189–91. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
9. Vander Mijnsbrugge W, Laleman W, Van Steenberghe W, Heye S, Verslype C, Maleux G. Long-term clinical and radiological outcome of endovascular embolization of pancreatitis-related pseudoaneurysms. *Acta Radiol.* 2017; 58: 316–22. [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

10. Kim J, Shin JH, Yoon HK *et al* Endovascular intervention for management of pancreatitis-related bleeding: a retrospective analysis of thirty-seven patients at a single institution. *Diagn. Interv. Radiol.* 2015; 21: 140–7. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
11. Flati G, Andren-Sandberg A, La Pinta M, Porowska B, Carboni M. Potentially fatal bleeding in acute pancreatitis: pathophysiology, prevention, and treatment. *Pancreas.* 2003; 26: 8–14. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
12. Tsiotos GG, Munoz Juarez MM, Sarr MG. Intraabdominal hemorrhage complicating surgical management of necrotizing pancreatitis. *Pancreas.* 1996; 12: 126–30. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
13. Zyromski NJ, Vieira C, Stecker M *et al* Improved outcomes in postoperative and pancreatitis-related visceral pseudoaneurysms. *J. Gastrointest. Surg.* 2007; 11: 50–5. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
14. Hyare H, Desigan S, Nicholl H, Guiney MJ, Brookes JA, Lees WR. Multi-section CT angiography compared with digital subtraction angiography in diagnosing major arterial hemorrhage in inflammatory pancreatic disease. *Eur. J. Radiol.* 2006; 59: 295–300. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
15. Hsu JT, Yeh CN, Hung CF *et al* Management and outcome of bleeding pseudoaneurysm associated with chronic pancreatitis. *BMC Gastroenterol.* 2006; 6: 3. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
16. Bhasin DK, Rana SS, Sharma V *et al* Non-surgical management of pancreatic pseudocysts associated with arterial pseudoaneurysm. *Pancreatol.* 2013; 13: 250–3. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
17. Rana SS, Kumar A, Lal A *et al* Safety and efficacy of angioembolisation followed by endoscopic ultrasound guided transmural drainage for pancreatic fluid collections associated with arterial pseudoaneurysm. *Pancreatol.* 2017; 17: 658–62. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
18. Hyare H, Desigan S, Brookes JA, Guiney MJ, Lees WR. Endovascular management of major arterial hemorrhage as a complication of inflammatory pancreatic disease. *J. Vasc. Interv. Radiol.* 2007; 18: 591–6. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
19. Okuno A, Miyazaki M, Ito H *et al* Nonsurgical management of ruptured pseudoaneurysm in patients with hepatobiliary pancreatic diseases. *Am. J. Gastroenterol.* 2001; 96: 1067–71. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
20. Marone EM, Mascia D, Kahlberg A, Brioschi C, Tshomba Y, Chiesa R. Is open repair still the gold standard in visceral artery aneurysm management? *Ann. Vasc. Surg.* 2011; 25: 936–46. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
21. De Rosa A, Gomez D, Pollock JG *et al* The radiological management of pseudoaneurysms complicating pancreatitis. *JOP.* 2012; 13: 660–6. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
22. Barge JU, Lopera JE. Vascular complications of pancreatitis: role of interventional therapy. *Korean J. Radiol.* 2012; 13(Suppl 1): S45–55. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
23. Golzarian J, Nicaise N, Deviere J *et al* Transcatheter embolization of pseudoaneurysms complicating pancreatitis. *Cardiovasc. Intervent. Radiol.* 1997; 20: 435–40. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
24. De Ronde T, Van Beers B, de Canniere L, Trigaux JP, Melange M. Thrombosis of splenic artery pseudoaneurysm complicating pancreatitis. *Gut.* 1993; 34: 1271–3. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
25. Vanlangenhove P, Defreyne L, Kunnen M. Spontaneous thrombosis of a pseudoaneurysm complicating pancreatitis. *Abdom. Imaging.* 1999; 24: 491–3. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
26. Sethi H, Peddu P, Prachalias A *et al* Selective embolization for bleeding visceral artery pseudoaneurysms in patients with pancreatitis. *Hepatobiliary Pancreat. Dis. Int.* 2010; 9: 634–8. [[PubMed](#)] [[Google Scholar](#)]