



Original Article

Comparison of Effects of Resistance Exercise and Endurance Exercise on Body Weight and Body Mass Index (Bmi)


Dr Shivam Bishnoi¹, Dr Sharda Singh², Dr Ankit Kumar Tiwari³, Dr Hariom Kumar Singh⁴

¹Assistant Professor, Department of Physiology, Autonomous State Medical College, Kaushambi, Uttar Pradesh

²Associate Professor, Department of Physiology, Autonomous State Medical College, Kaushambi, Uttar Pradesh

³Assistant Professor, Department of Biochemistry, Autonomous State Medical College, Kaushambi, Uttar Pradesh

⁴Professor, Department of Pharmacology, and Principal, Autonomous State Medical College, Kaushambi, Uttar Pradesh

 OPEN ACCESS

Corresponding Author:

Dr Shivam Bishnoi

Assistant Professor, Department of Physiology, Autonomous State Medical College, Kaushambi, Uttar Pradesh;

Email:

drshivambishnoi91@gmail.com

Received: 17-05-2026

Accepted: 08-06-2026

Available online: 18-06-2026

Copyright © International Journal of
Medical and Pharmaceutical Research

ABSTRACT

Background: Regular endurance exercise decreases obesity, improves BMI, reduces the level of inflammatory markers and chronic diseases; while the regular resistance exercise prevents the development of chronic diseases such as stroke, heart failure and other cardiovascular diseases.

Objective: The present study was designed to compare the effects of resistance and endurance exercise training on body weight and BMI of medical students at our institution.

Materials and Methods: This study was performed on male subjects who were apparently healthy with normal daily physical activity. There were sixty male participants, aged between 18-25yrs with normal BMI (18.5-24.9 kg/m²). Subjects were grouped into endurance (n=30) and resistance training (n=30). Endurance training and Resistance training was given for five days a week at 60-70% of maximum heart rate (HR) for 8 weeks. HR was measured before and after the exercise.

Results: The findings of the study indicate statistically significant differences in body weight & BMI before and after Endurance training as well as Resistance training. There is statistically significant decrease in body weight and BMI after both types of exercise. But endurance exercise training was found more effective in decreasing body weight and BMI.

Conclusion: Endurance exercise training is more effective in decreasing body weight and BMI.

Keywords: Endurance exercise; Heart rate; Resistance training; Body weight; BMI.

INTRODUCTION

Physical exercise in any form is necessary for everyone to be healthy and be free from diseases. Chronic diseases are result of reduced physical activity. Obesity has been related to development of chronic inflammation and chronic diseases such as diabetes, hypertension and myocardial infarction etc.^[1,2] Obesity is related to high BMI which is related to diabetes mellitus, coronary artery disease, hypertension and metabolic syndrome.^[3]

Regular endurance exercise decreases obesity, improves BMI and body fat composition.^[4,5] Regular endurance exercise reduces the level of inflammatory markers and chronic diseases.^[6] Regular resistance exercise prevents the development of chronic diseases such as stroke, heart failure and other cardiovascular diseases.^[7] Therefore the aims of present study were first to find out and second to compare, the effects of resistance and endurance exercise on body weight and BMI.

OBJECTIVE

The present study was designed to compare the effects of resistance and endurance exercise training on body weight and BMI of medical students at our institution.

MATERIALS AND METHODS

The study was performed at Autonomous State Medical College, Kaushambi, UP. Sixty male subjects were enrolled from the MBBS students studying in the college and their age group was between 18-25 years. Subjects selected were apparently healthy with normal daily physical activity and belong to normal BMI (18.5-24.9 kg/m²).^[8] A written informed consent was taken by subjects on prescribed consent form. Institutional ethical committee approved the ethical clearance for study. Two groups were created depending on the type of exercise, Group I - Resistance exercise training and Group II - endurance exercise training. Subjects were equally divided in each group (n=30). Subjects were performing exercise of moderate intensity (based on maximum heart rate) for 30 minutes daily. The Resistance group performed push-up, pull-up, and squats whereas the Endurance group performed jogging. Subjects were involved in the exercise training five days a week for eight weeks. In moderate intensity of exercise target heart rate was 64-76% of maximum heart rate. Heart rate assessment was done before the start of exercise and after five minutes of exercise when it reached 64-76% of maximum heart rate. It was measured by help of pulse oximeter. Follow-up was done after eight weeks of exercise in both the groups. Assessment of training program was done by comparison of changes in body weight and BMI. **Body mass index (BMI)** is calculated as the body mass, in kilograms (kg), divided by the square of the body height, in square meters (m²).^[9]

Classification of intensity of exercise based on maximum heart rate^[10]

Intensity of exercise	HRmax%
Very light	57
Light	57-63
Moderate	64-76
Vigorous	77-95
Maximal	96-100

According to American college of sports medicine guidelines 2018

Heart rate maximum:

HRmax = 220 – age in years (beats per minutes)^[11]

Statistical analysis: The data was assessed by using Statistical Package for Social Science (SPSS) version 21.0. All qualitative variables were mentioned in form of frequency (%). All the quantitative variables were mentioned in form of Mean±SD. The data was assessed using student t-test and paired t-test. Student t-test was used for comparing body weight and BMI between the two groups. Paired t-test was used for intragroup comparison. The p value less than 0.05 was considered significant.

RESULTS

Mean age of subjects of group I was found to be higher (19.98±1.26 years) as compared to group II (19.97±0.98 years). Mean pre intervention BMI of both the groups was found to be similar (21.97±1.78 kg/m²). (Table-1)

Range of pre-intervention body weight in both the groups was 54-83 kg, though pre-intervention body weight of subjects in Group I (66.33±9.59 kg) was found to be higher as compared to Group II (65.87±9.47 kg) but this difference was not found to be significant statistically (p=0.850). Post-intervention body weight of overall population was 53-83 kg, body weight of Group I (65.67±9.23 kg) was still higher than that of Group II (63.53±9.10 kg) and difference in post-intervention body weight between two groups was not significant statistically. (Table-2).

After intervention statistically significant decline in pre-intervention body weight was observed in both the groups. In Group I decline of 0.67±0.80 kg was observed while in Group II decline of 2.33±0.80 kg was observed. Percentage post-intervention change in body weight was 1.01% in Group I and 3.54% in Group II. Group II (Endurance exercise) was more effective for losing body weight. (Table-3)

Range of pre-intervention BMI in overall as well as both the groups was 18.60 to 24.90 kg/m². Mean pre-intervention BMI of both the groups was found to be similar (21.97±1.78 kg/m²). Range of post-intervention BMI in overall study population and in Group I was 18.50 to 24.60 kg/m² while same in Group II was 18.50 to 23.60 kg/m². Mean post-intervention BMI of Group I (21.49±1.51 kg/m²) was found to be higher than that of Group II (20.66±1.42 kg/m²). This difference was found to be significant statistically. (Table-4).

After intervention significant decline in pre-intervention body mass index was observed in both the groups. In Group I decline in BMI was 0.47±0.70 kg/m² while in Group II decline in BMI was 1.31±0.67. On evaluating the % post-intervention change in Group I (2.15%) and Group II (5.95%), it was observed that decline in BMI in Group II was higher as compared to Group I. (Table-5)

Table-1: Socio-demographic characteristics

Parameter	Resistance exercise	Endurance exercise
Age (Mean±SD) years	19.98±1.26	19.97±0.98

Weight (Mean±SD) kg	66.33±9.59	65.87± 9.47
BMI (Mean±SD) kg/m²	21.97±1.78	21.97±1.78
Race	Asian	Asian
Type of exercise	Push-ups, pull-ups, squats	Jogging

Table 2: Between Group Comparison of Pre and Post-intervention Body Weight (in kg)

Group	Range		Mean	S.D.	95% CI	
	Min.	Max.			Lower bound	Upper bound
Pre-intervention						
Group I (n=30)	54.00	83.00	66.33	9.59	62.75	69.91
Group II (n=30)	54.00	83.00	65.87	9.47	62.33	69.40
Total (n=60)	54.00	83.00	66.10	9.45	63.66	68.54
Student 't' test			't'=0.190; p=0.850 (NS)			
Post-intervention						
Group I (n=30)	55.00	83.00	65.67	9.23	62.22	69.11
Group II (n=30)	53.00	81.00	63.53	9.10	60.14	66.93
Total (n=60)	53.00	83.00	64.60	9.15	62.24	66.96
Student 't' test			't'=0.902; p=0.371 (NS)			

Table 3: Intragroup Change in Pre and Post-intervention Body Weight (Paired 't' test)

Group	Mean change	S.D.	% Post-intervention Change	't'	'p'
Group I	-0.67	0.80	-1.01	-4.551	<0.001
Group II	-2.33	0.80	-3.54	-15.930	<0.001
Total	-1.50	1.16	-2.27	-10.041	<0.001

Table 4: Between Group Comparison of Pre and Post-intervention Body Mass Index (in kg/m²)

Group	Range		Mean	S.D.	95% CI	
	Min.	Max.			Lower bound	Upper bound
Pre-intervention						
Group I (n=30)	18.60	24.90	21.97	1.78	21.30	22.63
Group II (n=30)	18.60	24.90	21.97	1.78	21.30	22.63
Total (n=60)	18.60	24.90	21.97	1.77	21.51	22.42
Student 't' test			't'=<0.001 (Sig); p=0.1.000(NS)			
Post-intervention						
Group I (n=30)	18.50	24.60	21.49	1.51	20.93	22.06
Group II (n=30)	18.50	23.60	20.66	1.42	20.13	21.19
Total (n=60)	18.50	24.60	21.08	1.52	20.69	21.47
Student 't' test			't'=2.198; p=0.032 (Sig)			

Table 5: Intragroup Change in Pre- and Post-intervention Body Mass Index (Paired 't' test)

Group	Mean change	S.D.	% Post-intervention Change	't'	'p'
Group I	-0.47	0.70	-2.15	-3.708	0.001
Group II	-1.31	0.67	-5.95	-10.748	<0.001
Total	-0.89	0.80	-4.05	-8.653	<0.001

DISCUSSION-

Subjects who completed resistance exercise training also showed statistically significant decrease in body weight and BMI after exercise. Our findings are similar to Donges et al,^[12] Chun-De Liao,^[13] and Hee-Jae KIM et al,^[14] who studied the effect of resistance exercise on body fat composition and observed that there is statistically significant decrease in body weight and BMI after exercise.

Subjects who performed endurance exercise training showed statistically significant decrease in body weight and BMI. There was statistically significant increase in lean mass after exercise. These findings are in accordance to Sayyed Mohammad Marandi et al,^[15] Hee Jae KIM et al,^[14] J. K. Song et al,^[16] and Tugay Yilmaz et al,^[17] who studied the effect of endurance exercise on body fat composition and observed that there is statistically significant decrease in body weight

and BMI after exercise. In the present study positive correlation of endurance and resistance exercise was found on body weight and BMI. But endurance exercise training was more effective in improving body weight and BMI.

CONCLUSION

There is statistically significant decrease in weight and BMI after resistance exercise training as well as after endurance exercise training. However, the decrease in weight and BMI is more effective in endurance exercise group than in resistance group.

ACKNOWLEDGEMENT

The authors thank the MBBS students of Autonomous State Medical College, Kaushambi, UP for their active participation. The authors also wish to acknowledge Dr. Shailendra Vashistha (Assistant Professor, Transplant Immunology-HLA Lab, Dept of IHTM, GMC, Kota) and the VAssist Research team (www.thevassist.com) for their contribution in manuscript editing and submission process.

Conflict of Interest: None

Source of Funding: Nil

REFERENCES

1. Ellulu MS, Patimah I, Abed Y. Obesity and inflammation: The linking mechanism and the complications. *Arch Med Sci.* 2016 Mar;13(4):851–63. Doi: [10.5114/aoms.2016.58928](https://doi.org/10.5114/aoms.2016.58928). PMID: 28721154 Published online 2017.
2. Corbin KD. Obesity in type 1 diabetes: Pathophysiology, Clinical Impact, and Mechanisms. *Endocr Rev.* 2018 Oct;39(5):629-63. Doi: [10.1210/er.2017-00191](https://doi.org/10.1210/er.2017-00191) PMID: 30060120
3. Kearns K, Dee A, Fitzgerald AP, Doherty E, Perry IJ. Chronic disease burden associated with overweight and obesity in Ireland: The effects of a small BMI reduction at population level. *BMC Public Health.* 2014 Feb;14:143-9. Doi: [10.1186/1471-2458-14-143](https://doi.org/10.1186/1471-2458-14-143). PMID: 24512151, 2014.
4. Durstine JL, Haskell WL. Effects of exercise training on plasma lipids and lipoproteins. *Exerc Sport Sci Rev.* 1994; 22:477- 521.
5. Chi CH, Ko MC, Wu LS, Yeh DP, Kan NW, Lee PF, et al. Benefits of different intensity of aerobic exercise in modulating body composition among obese young adults: A pilot randomized controlled trial. *Health Qual Life Outcomes.* 2017 Aug;15(1):168-75. Doi: [10.1186/s12955-017-0743-4](https://doi.org/10.1186/s12955-017-0743-4)
6. Zheng G, Qiu P, Xia R, Lin H, Ye B, Tao J, et al. Effect of aerobic exercise on inflammatory markers in healthy middle-aged and older adults: A systematic review and meta-analysis of randomized controlled trials. *Front Aging Neurosci.* 2019 Apr;11:98-105. Doi: [10.3389/fnagi.2019.0098](https://doi.org/10.3389/fnagi.2019.0098)
7. Braith RW, Stewart KJ. Resistance exercise training: its role in the prevention of cardiovascular disease. *Circulat.* 2006 Jun; 113 (22) : 2642 – 50. doi: [10.1161/CIRCULATIONAHA.105.584060](https://doi.org/10.1161/CIRCULATIONAHA.105.584060)
8. The SuRF Report 2. The Surveillance of Risk Factors Report Series (SuRF). World Health Organization. 2005. p. 22. Archived (PDF) from the original on 2022-10-09.
9. Chaudhary S, Kang MK, Sandhu JS. The effects of aerobic versus resistance training on cardiovascular fitness in obese sedentary females. *Asian J Sports Med.* 2010 Dec;1(4):177-84. Doi: [10.5812/asjms.34835](https://doi.org/10.5812/asjms.34835). PMID: 22375205, PMID: PMC3289184.
10. American college of sports medicine (2018). ACSMs guidelines for exercise testing and prescription 10th edition. Philadelphia: Wolters kluwer.
11. Almaadawy O, Uretsky BF, Krittanawong C, Birnbaum YJ. Target heart rate formulas for exercise stress testing: What is the evidence? *Clin Med.* 2024 Sep;13(18):5562. doi: [10.3390/jcm13185562](https://doi.org/10.3390/jcm13185562).
12. Donges. Effects of resistance or aerobic exercise training on interleukin-6, c-reactive protein, and body composition, *medicine & science in sports & exercise.* 2010 Feb;42(2):304-13.
13. Liao CD. Effects of elastic resistance exercise on body composition and physical capacity in older women with sarcopenic obesity. *Medicine (Baltimore).* 2017 Jun;96(23):e7115. Doi: [10.1097/MD.00000000000007115](https://doi.org/10.1097/MD.00000000000007115). PMID: PMC5466239, PMID: 28591061.
14. Kim HJ. Effect of aerobic training and resistance training on circulating irisin level and their association with change of body composition in overweight/obese adults: A pilot study. *Physiol Res.* 2016Jun;65(2):2719-26. Doi:[10.33549/physiolres.93299](https://doi.org/10.33549/physiolres.93299).
15. Marandi SM, Abadi NGB, Ghasemi G. Effects of intensity of aerobics on body composition and blood lipid profile in obese/overweight females. *Int J Prev Med.* 2013 Apr;4(Suppl 1):S118-25. PMID: 23717761. PMID: PMC3665017.
16. Song JK. Effects of 12 weeks of aerobic exercise on body composition and vascular compliance in obese boys. *J Sports Med Phys Fitness.* 2012 Oct;52(5):522-9. PMID: 22976739.
17. Yilmaz T, Daglioglu O. The effect of aerobic training program on cardiopulmonary parameters and oxygen saturation in elite Judokas. *Turkish J Sport Exerc.* 2018;20(3):333–7. Doi: [10.15314/tsed.471231](https://doi.org/10.15314/tsed.471231).