



Original Article

Prevalence of Malnutrition in Beta Thalassemia Major Patients in a Government Tertiary Care Centre in South India

Ashwini G Rayee^{1*}, Sharanabasappa S Dhanwadkar², Babalala Kadegaon³, Revanasiddappa Bhosgi⁴, Ruthu G Rayee⁵

¹Senior resident, Department of Paediatrics, Adichunchanagiri institute of medical sciences, B G nagara, Mandya district, Karnataka, India

²Associate professor, Department of Paediatrics, Gulbarga Institute of Medical Sciences, Kalaburagi, Karnataka, India

³Assistant professor, Department of Paediatrics, Gulbarga Institute of Medical Sciences, Kalaburagi, Karnataka, India

⁴Intern, Adichunchanagiri institute of medical sciences, B G nagara, Mandya district, Karnataka, India

⁵Professor and HOD, Department of Paediatrics, Gulbarga Institute of Medical Sciences, Kalaburagi, Karnataka, India

OPEN ACCESS

Corresponding Author:

Dr. Ashwini G Rayee

Senior resident Department of
Paediatrics, Adichunchanagiri
institute of medical sciences, B G
nagara, Mandya district,
Karnataka, India

Email: rayeeashwinig@gmail.com

Received: 21-06-2026

Accepted: 10-07-2026

Available online: 25-07-2026

Copyright © International Journal of
Medical and Pharmaceutical Research

ABSTRACT

Background and Objectives: Beta thalassemia is a hereditary hemoglobinopathy characterized by the defects in the beta globin chain of haemoglobin. In India, For about 12000 infants born every year have major form of the disease and half of these patients die before reaching adulthood. Large proportion of these early deaths are due to malnutrition. With this background, this study was conducted with the objective to know the prevalence of malnutrition in Beta thalassemia major patients.

Methods: The present cross sectional observational study was carried among 100 children aged 6 months to 18 years with confirmed beta thalassemia major patients. A detailed history taking, examination was done. Based on BMI, malnutrition was assessed using age and sex specific WHO BMI charts. Statistical data was analysed by IBM SPSS 25.0 version software and statistical significance for all analyses was set at $P < 0.05$.

Results: In our study, about 36.0% children were belonging to the age group of 0.5—5 years and 64.0% children were belonging to the age group of 15.1-18.0 years.

In this age group of 6 months to 5 years 11.1% children were malnourished, In this age group of 5.1 to 18 years, malnourished children were 71.9%. Among the study participants, 50.0% were malnourished with a mean body mass index of 14.46 kg/m².

Interpretation & Conclusion: In our study half of the study participants had malnutrition, this shows that malnutrition is also one of the entity which should be taken care in these patients.

Keywords: Malnutrition; Thalassemia major; Body Mass Index.

INTRODUCTION

Thalassemia is autosomal recessive single gene disorder. About 200 million people globally are affected by thalassemia, with half of these cases having the severe form of the disease that is beta thalassemia major. ¹⁻³

Beta thalassemia is a hereditary hemoglobinopathy characterized by the defects in the beta globin chain of haemoglobin. Homozygous and compound heterozygous forms have an imbalance in the production of alpha and non alpha globin chains, which results in ineffective erythropoiesis and decreased production of normal haemoglobin A. ⁴

In India, For about 12000 infants born every year have major form of the disease {10% of global burden} and half of these patients die before reaching adulthood. Large proportion of these early deaths are due to malnutrition. ⁵

There is multi factorial association for malnutrition in thalassemia. According to the existing literature, children with thalassemia have an average energy intake lower than the recommended daily dietary allowances. Most of them belong to a more economically weaker section of the society. As a result, thalassemia patients develop multiple micronutrients deficiency which can be quoted as “hidden hunger”. So, socio economic status and literacy of parents play a major role in this regard.⁵

So, this study was taken up to assess the malnutrition status in beta thalassemia patients, So that early identification of malnutrition in these patients will help in the early management of malnutrition, so that the quality of life of these patients can be improved and thereby helping to reduce the early deaths associated with malnutrition in thalassemia major patients.

METHODOLOGY:

This cross sectional observational study was conducted from August 2022 to January 2025 among 100 confirmed cases of beta thalassemia major children attending department of Paediatrics, Gulbarga institute of Medical sciences, Kalaburagi in the age group of 6months to 18years. Beta thalassemia major children who are having neurological disorders, complex syndromes/genetic disorders, chronic kidney disease or chronic lung disease were excluded from the study.

The study was conducted after obtaining the ethics committee approval from Institutional Ethics committee. Informed consent was obtained from all eligible participants after explaining the objectives and nature of this study in their own language.

Malnutrition was assessed in the study subjects as soon as they are first seen and enrolled into the study. A detailed history taking, examination and relevant investigations was done. Details such as age, sex, place of residence (rural or urban), caste, per capita monthly income , socio economic status (classified based on modified B G Prasad classification)⁶, pre transfusion haemoglobin level (gm/dl), number of blood transfusion received last year, splenectomy status, history of intake of iron chelators , duration since receiving iron chelators were taken. On examination, we looked for presence/absence of thalassemia facies. Spleen size and liver size were measured. Length (for children less than 2 years), Height (for children more than 2 years), Weight was measured and BMI was calculated. Based on BMI, malnutrition was assessed using WHO BMI charts^{7,8} and the effect of above factors on the malnutrition was assessed.

In our study, malnutrition was assessed in beta thalassemia major patients using WHO BMI growth charts. The BMI was plotted on age and sex specific WHO BMI charts. WHO BMI charts of birth to 5 years was used for children aged 6 months to 5 years.⁷ And WHO BMI charts of 5 to 19 years was used for the children aged 5 to 18 years.⁸

The nutritional status of 6 months to 5 years old children was classified according to BMI based WHO classification for birth to 5 years of age as follows: ⁹

- Obese: BMI for age greater than +3 standard deviations (SD) of the median.
- Overweight: BMI for age greater than +2 SD but equal or less than +3 SD of the median.
- Normal: BMI for age equal or less than +2 SD and but more than –2 SD of the median.
- Moderate acute malnutrition: BMI for age equal or less than –2 SD but equal or more than –3 SD of the median
- Severe acute malnutrition: BMI for age less than –3 SD of the median.
- The nutritional status of 5-18 years old children was classified according to BMI based WHO classification for 5-19 years of age as follows: ⁸
- Obese: BMI for age more than +2 SD.
- Overweight: BMI for age more than +1 SD but equal or less than +2 SD.
- Normal: BMI for age equal or more than -2 SD but equal or less than +1 SD.
- Thin: BMI for age less than -2 SD but equal or more than -3 SD.
- Severely thin: BMI for age less than –3 SD of the median.
- Statistical data was analyzed by using statistical package for social sciences (SPSS, version 20.0). If p value is <0.05, it is considered as statistically significant.

RESULTS:

Table 1: Age and gender wise distribution of children

Age in years	Males		Females		Total	
	Number	Percentage	Number	Percentage	Number	Percentage
0.5—5 years	20	33.9	16	39.0	36	36.0
5.1—18.0 years	39	66.1	25	60.9	64	64.0
Total	59	100.0	41	100.0	100	100.0
Mean ± SD	7.57 ± 4.48		8.05 ± 5.05		7.77 ± 4.68	

Table 1 shows that, 36 (36.0%) children were belonging to the age group of 0.5—5 years, 64(64.0%) children were belonging to the age group of 5.1—18.0 years. Minimum age of children in the study was 1.5 years and maximum age was 17.25 years.

The mean age of male children was 7.57 years, and the mean age of female children was 8.05 years. Mean age of all children was 7.77 years. In the study, Male children were 59 (59.0%) and female children were 41 (41.0%).




Figure 1: Age and Gender wise distribution of Children

In the study, 65 (65.0%) children were residing in rural area and 35 (35.0%) children were residing in urban area.

In the present study, 91 (91.0%) children were belonging to Hindu Religion and 9 (9.0%) children were belonging to Muslim. Among Hindu religion, 51 (51.0%) children were belonging to SC/ST (scheduled caste/scheduled tribe), followed by 34 (34.0%) children were belonging to OBC category .

Table 2: Socio-economic status wise distribution of children

Socio economic status Classification	Number of children	Percentage (%)
Upper Class	0	0.0
Upper Middle Class	4	4.0
Middle Class	23	23.0
Lower Middle Class	61	61.0
Lower Class	12	12.0
Total	100	100.0

Table 2 shows, Socio-economic status wise distribution of children (As per modified B G Prasad classification). Majority of children, 61 (61.0%) were belonging to Lower middle class, followed by 23 (23.0%) children were belonging to middle class, 12 (12.0%) children were belonging to lower class, 4 (4.0%) children were belonging to upper middle class and none of the children were belonging to upper class.

In our study, it was observed that; the mean height of children was 103.16 cm, the mean weight of children was 15.83 kg, the mean BMI of children was 14.46

Table 3: Age wise classification of BMI and prevalence of malnutrition in beta thalassemia major children

Age	BMI categories	Number of children	Percentage
6 months to ≤ 5 years	SAM	0	0.0
	MAM	4	11.1
	Normal	32	88.9
	Over weight	0	0.0
	Obese	0	0.0
Sub total	Total	36	100.0
5.1—18.0 years	Severe Thin	28	43.8
	Thin	18	28.1

	Normal	17	26.6
	Over weight	1	1.5
	Obese	0	0.0
Sub total	Total	64	100.0
Grand Total	---	100	100.0

Table 3 shows: In the study, out of 36 beta thalassemia major children aged 6 months to 5 years, 4 (11.1%) children were malnourished and were belonging to BMI category of MAM and 32 (88.9%) children were belonging to normal BMI category and None of the Children were seen in the BMI categories of SAM, Overweight and Obese.

Study observed that 64 (64.0%) children were belonging to the age of 5.1 to 18 years, among them 28 (43.8%) children were belonging to BMI category of severely thin, 18 (28.1%) children were belonging to BMI category of thin, 17 (26.6%) children were belonging to BMI category of normal and only 1 (1.5%) child was overweight and None of the children were seen in the obese category of BMI. In this age group of 5.1 to 18 years, malnourished children were 46 (71.9%). So in the study the prevalence of malnutrition in beta thalassemia major children in the age group of 6 months to 18 years was 50.0%. Total number of malnourished children in the study was 50. The children in the age group of 5.1 to 18 years had higher percentage of malnourished children as compared to the children in the age group of 6 months to 5 years.




Figure 2: Age wise classification of BMI and prevalence of malnutrition in beta thalassemia major children.

Table 4: Gender wise classification of BMI and prevalence of malnutrition in beta thalassemia major children

Age	Gender	BMI categories	Number of children	Percentage (%)
6 months to ≤ 5 years	Male (N=20)	MAM	2	5.6
		Normal	18	50.0
	Female (N=16)	MAM	2	5.6
		Normal	14	38.8
Sub total	---	Total	36	100.0
5.1—18.0 years	Male (N=39)	Severe Thin	16	25.0
		Thin	9	14.0
		Normal	13	20.3
		Over weight	1	1.6
	Female (N=25)	Severe Thin	12	18.8
		Thin	9	14.0
		Normal	4	6.3
		Over weight	0	0.0
Sub total	----	Total	64	100.0
Grand Total	----	---	100	100.0

Table 4 shows that, in the study, 36 children were in the age group of 6 months to 5 years of age; Among them 20 (55.5%) were male children and 16 (44.4%) were female children. Malnutrition were observed in 4 (11.1%) children in this age

group, among them 2 (50.0%) were males and 2(50.0%) were females. The percentage of malnourished children is same in both males and females belonging to the age group of 6 months to 5 years.

Out of 100 study children; 64 (64.0%) of children were in the age group of 5.1—18.0 years. Among them 39 (60.9%) were male children and 25 (39.1%) were female children. Among male children, 25 (64.1%) were malnourished and among female children 21 (84.0%) were malnourished. The percentage of malnourished children is higher among female gender as compared to males belonging to the age group of 5.1 to 18 years.

DISCUSSION:

Thalassemia is autosomal recessive single gene disorder. About 200 million people globally are affected by thalassemia, with half of these cases having the severe form of the disease that is beta thalassemia major.¹⁻³

In India, For about 12000 infants born every year have major form of the disease {10% of global burden} and half of these patients die before reaching adulthood. Large proportion of these early deaths are due to malnutrition.⁵

There is multi factorial association for malnutrition in thalassemia. According to the existing literature, children with thalassemia have an average energy intake lower than the recommended daily dietary allowances. Most of them belong to a more economically weaker section of the society. As a result, thalassemia patients develop multiple micronutrients deficiency which can be quoted as “hidden hunger”.⁵

So, this study was taken up to assess the malnutrition and various factors influencing malnutrition in beta thalassemia patients, So that early identification of malnutrition in these patients will help in the early management of malnutrition. The present cross sectional observational study was carried out among 100 children with beta thalassemia major in the Department of Pediatrics, Gulbarga Institute Of Medical Sciences, Kalaburagi, Karnataka to know the prevalence of malnutrition in Beta thalassemia major patients

Table 5: Sample size comparison with various studies

	Present study	Bijit Biswas et al. ⁵	Mirhosseini et al. ¹⁰	Upadhye et al. ¹¹	Fahim et al. ¹²
Sample size	100	328	140	122	100

The sample size in the present study is comparable to the other studies as mentioned in the Table 5 except for study conducted by Bijit Biswas et al,⁵ where the sample size was larger compared to our study.

Socio demographic profile of the participants in the study

- In our study, about 36.0% children were belonging to the age group of 0.5—5 years and 64.0% children were belonging to the age group of 15.1-18.0 years.
- Minimum age of children in the study was 1.5 years and maximum age was 17.25 years.
- The mean age of male children was 7.57 years, and the mean age of female children was 8.05 years. Mean age of all children was 7.77 years.
- In our study, Male children were 59.0% and female children were 41.0%.
- In a study conducted by Bijit Biswas et al,⁵ Most of the study participants (37.2%) were aged between 11 and 12 years (range, 5–12 years). The mean age in the study was 8.0 years. There was almost equal representation of both sexes that is 54.0% were males and 46.0% were females.
- The study conducted by Mirhosseini et al,¹⁰ had participants aged 8-18 years. The mean age in the study was 13.5 years and this study included 56.4 % boys and 43.6 % girls. In a study conducted by Sheikh et al.¹³ age range of the patients of thalassemia major was 2 to 16 years with mean age of 7.88 years.
- In our study, 65.0% children were residing in rural area and 35.0% children were residing in urban area and majority (51.0%) of the children were belonging to SC/ST (scheduled caste/scheduled tribe). In a study conducted by Bijit Biswas et al, ⁵ 51.1% children were residing in rural area and 40.7% children were residing in urban area.
- In our study, Majority of children, 61.0% were belonging to Lower middle class, followed by 23.0% children were belonging to middle class. Similar observations were seen in a study conducted by Bijit Biswas et al, ⁵ where 48.8% were belonging to lower middle class followed by 24.1% were belonging to middle class.

Age wise classification of BMI and prevalence of malnutrition in beta thalassemia major children

In our study, out of 36 beta thalassemia major children aged 6 months to 5 years, 11.1% children were malnourished and were belonging to BMI category of MAM and 88.9% children were belonging to normal BMI category and None of the Children were seen in the BMI categories of SAM, Overweight and Obese. 64.0% children were belonging to the age of 5.1 to 18 years, among them 43.8% children were belonging to BMI category of severely thin, 28.1% children were

belonging to BMI category of thin, 26.6% children were belonging to BMI category of normal and only 1.5% child was overweight and None of the children were seen in the obese category of BMI. In this age group of 5.1 to 18 years, malnourished children were 71.9%. So in our study the prevalence of malnutrition in beta thalassemia major children in the age group of 6 months to 18 years was 50.0%. Total number of malnourished children in the study was 50. The children in the age group of 5.1 to 18 years had higher percentage of malnourished children as compared to the children in the age group of 6 months to 5 years.

The mean BMI of children in our study was 14.46. Similar to our study, In the study conducted by Bijit Biswas et al,⁵ mean BMI was 13.9 and in the study conducted by Sheikh et al,¹³ mean BMI was 14.03. Where as slightly higher mean BMI was reported in a study conducted by Mirhosseini et al,¹⁰ which is about 17.6.

In the present study, half (50.0%) of the study participants were malnourished, which was higher than the value reported in a study of Pemde et al,¹⁴(24.2%), study of Mirhosseini et al,¹⁰(33.6%) and a study of Trehan et al,¹⁵ (26.7%) and similar to the value reported in a study of Qaisar I et al,¹⁶ (50.6%) study of Sheikh et al,¹³ (58.69%) ,study of Bijit Biswas et al,⁵ (48.2%) and a study of Fahim et al,¹² (47.0%).The variability of the findings may be attributed to the differences in the participants' age, ethnicity, and geographical plausibility.

In our study, BMI was plotted on age and sex specific WHO growth charts and other studies like Qaisar I et al,¹⁶ and Pemde et al,¹⁴ also have used WHO growth charts.

Gender wise classification of BMI and prevalence of malnutrition in beta thalassemia major children

In our study, Malnutrition was observed in 11.1% children in the age group of 6 months to 5 years, among them 50.0% were males and 50.0% were females. The percentage of malnourished children is same in both males and females belonging to the age group of 6 months to 5 years. In age group of 5.1 to 18 years, 64.1% male children were malnourished and 84.0% female children were malnourished. The percentage of malnourished children is higher among female gender as compared to males belonging to the age group of 5.1 to 18 years. In a study conducted by Mirhosseini et al,¹⁰ prevalence of malnutrition was 44.3 % for boys and 19.6 % for girls. In a study conducted by Bijit Biswas et al,⁵ prevalence of malnutrition was 49.0% among females and 47.5% among males. In a study conducted by Qaisar I et al,¹⁶ prevalence of malnutrition was 34.4% among females, 31.6% among males in the age group of 0 to 59 months and prevalence of malnutrition was 66.6% among females and 65.8% among males in the age group of 60 to 215 months in the same study.so in all the above studies prevalence of malnutrition was more in females when compared to males which is similar to our study except for study conducted by Mirhosseini et al,¹⁰ where prevalence of malnutrition is more seen in boys.

CONCLUSION:

In our study the prevalence of malnutrition in beta thalassemia major children in the age group of 6months to 18 years was 50.0%. So periodic growth assessment once in every 3 months is necessary for early detection and management of malnutrition in these patients for better outcome.

The children with older age group of 5.1 to 18 years were at higher risk of malnutrition. indicating that as disease progresses the risk of malnutrition increases attributed to the disease progression as well as hidden hunger in these patients because of increased calorie requirements.

ACKNOWLEDGEMENT

Authors acknowledge the help received from scholars whose articles have been cited and included in references of this manuscript.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee.

REFERENCES

1. Kasper DL, Fauci AS, Hauser SL, Longo DL, Jameson JL, Loscalzo J. Harrison's principles of internal medicine. 19th ed. New York (NY): McGraw-Hill Medical Publishing; 2015.
2. Colah R, Gorakshakar A, Nadkarni A. Global burden, distribution and prevention of beta-thalassemias and hemoglobin E disorders. *Expert Rev Hematol* 2010; 3(1):103-117.
3. Sharma N, Gupta A, Gupta G. Assessment of serum ferritin, vitamin B12 and folic acid level in thalassemia. *J Med Sci Clin Res* 2017; 5(7):29910-29914.
4. Kattamis A, Forni GL, Aydinok Y, Viprakasit V. Changing patterns in the epidemiology of β -thalassemia. *Eur J Haematol* 2020; 105(6):692-703.
5. Biswas B, Naskar NN, Basu K, Dasgupta A, Basu R, Paul B. Malnutrition, its attributes and impact on quality of life: an epidemiological study among beta thalassemia major children. *Korean J Fam Med* 2021; 42(1):66-72.
6. Mangal A, Kumar V, Panesar S, Talwar R, Raut D, Singh S. Updated BG Prasad socioeconomic classification, 2014: A commentary. *Indian J Public Health* 2015; 59:42-44.

7. <https://www.who.int/toolkits/child-growth-standards/standards/body-mass-index-for-age-bmi-for-age> [cited 2023 Jan 18].
8. <https://www.who.int/tools/growth-reference-data-for-5to19-years/indicators/bmi-for-age> [cited 2023 Jan 18].
9. World Health Organization. Assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition: Updates for the integrated management of childhood illness (IMCI) – Guideline. Geneva: World Health Organization; 2017.
10. Mirhosseini NZ, Shehar S, Mobarhan MG, Kamaruddin NA, Banihasham A, Yusoff N, et al. Factors affecting nutritional status among pediatric patients with transfusion-dependent beta thalassemia. *Mediterr J Nutr Metab* 2013;6:45-51.
11. Upadhye M, Rajput U, Kinikar A. Prevalence of malnutrition in pediatric beta-thalassemia major patients admitted to a tertiary care center. *Global J Res Anal* 2018; 7:70-72.
12. Fahim FM, Saad K, Askar EA, Eldin EN, Thabet AF. Growth parameters and vitamin D status in children with thalassemia major in upper Egypt. *Int J Hematol Oncol Stem Cell Res* 2013; 7:10-14.
13. Sheikh MA, Shakir MU, Shah M. The assessment of nutritional status of children with beta thalassemia major with body mass index. *Pak J Med Health Sci.* 2017; 11:262-265.
14. Pemde HK, Chandra J, Gupta D, Singh V, Sharma R, Dutta AK. Physical growth in children with transfusion-dependent thalassemia. *Pediatr Health Med Ther* 2011; 2:13-19.
15. Trehan A, Sharma N, Das R, Bansal D, Marwaha RK. Clinicoinvestigational and demographic profile of children with thalassemia major. *Indian J Hematol Blood Transfus* 2015; 31:121-126.
16. Qaisar I, Rehman A, Jillani KS. Assessment of nutritional status of children with beta thalassemia major with BMI. *Med Forum* 2020; 31:107-110.