

**Growth Performance And Economic Impact Of Broilerchicken Fed Concentrates Supplemented With Processed Cocoa Pod Waste****R. P. Obongekpe\***

Department of Animal Science, University of Uyo, Uyo, Nigeria\*

**ABSTRACT**

The study was conducted to examine the performance growth and economic impacts of broiler chicken fed concentrates supplemented with processed cocoa pod waste. The experiment was conducted at the Teaching and Research Farm, University of Uyo, Uyo, Akwa Ibom State. Cocoa pod waste or husk was collected and processed by ash treatment to form a processed cocoa pod husk meal (PCHM). Three experimental diets were formulated at both the starter and finisher phases, in which PCHM was included at 0, 5 and 10% and designated as diets 1, 2 and 3, respectively. One hundred and eighty 1-day-old Arbor Acres broiler chicks were randomly distributed to three dietary treatments (10 birds/replicate; 60 birds/treatment) in a completely randomized design. The growth performance, carcass, relative internal organ weights and haemato biochemical indices were determined. Histological examination of the liver and heart samples was also determined. Results: The PCHM inclusion did affect ( $P < 0.05$ ) the performance characteristics of the broiler chicks and also for the feed intake that significantly ( $P < 0.05$ ) increased in birds fed 8% of PCHM-inclusive diet at the starter phase. The carcass traits, relative internal organ weights, haematological indices and serum biochemical indices of the broiler chickens were similar ( $P > 0.05$ ) across the dietary treatments. The serum glutathione peroxidase and catalase concentration were higher ( $P < 0.05$ ) in birds fed PCHM-inclusive diets compared to those fed the control diet. Implications and recommendations were made from the findings of the study.

**Keywords:** *Broilers, cocoa pod waste, Growth performance, Economic impacts.***\*Corresponding Author****R. P. Obongekpe**Department of Animal Science, University of Uyo,  
Uyo, Nigeria

© Copy Right, IJMPR, 2020. All Rights Reserved

**INTRODUCTION:**

Feeds and feeding constitutes a larger proportion of the expenditure in the poultry industry especially when ingredients like grains are used. According to Nworgu et al. [1] feeding cost covers between 60 to 70% of the total production cost of monogastric animals. Therefore, replacement of one or more of the major conventional feed ingredients with cheap and available non-conventional feed ingredients will have reducing effects on animal feed production cost [2]. The use of agro-wastes such as cassava peels and cocoa bean shell in monogastric animal production was reported [3]; [4], [5]; [6], [7] to have positive effects on monogastric animals especially poultry. Feeding of poultry with phytochemicals (products derived from plant, e.g. dried plant material, essential oil, pure isolated compound, or extract, which contain secondary plant metabolites) has been reported as a reliable means of combating the negative effect of oxidative stress in heat-stressed poultry [8].

Cocoa pod husk is a typical under-utilized agro-waste from the commercial cocoa farm, which could provide nutritional benefits to monogastric animal production [2]. Cocoa pod husk forms about 70% (w/w) of the whole mature cocoa fruit; has low crude protein (9.14%), high crude fibre (35.78%) [9] and anti-nutritional factors such as threobromine (2.64%); and has tannin (0.91%), caffeine (1.14%) and high fibre [2]. However, it was also reported that the optimal utilization of agro-waste in monogastric animal production is hampered by the anti-nutritional factors which cause the inactivation of some nutrients, diminution of metabolic utilization of food or the process of food digestion [10]. Therefore, various treatments such as fermentation [11]; [12], ash treatment [13], enzyme supplementation [7],[14], soaking and sun-drying Adebawale[1] as cited in [15] among others have been used to improve the nutritive value of agro-waste. In particular, combination of ash treatment with fermentation was also reported to improve the nutritive values of cocoa pod husk meal and its suitability in monogastric animal production [2]. Dietary inclusion of processed cocoa pod husk meal up to 150 g/kg was reported to support normal growth performance, carcass traits and relative internal organ weights in the rabbits [15].

Variation could occur in performance response of different species of animals to unconventional feed ingredients,

and presently, relatively few works had been reported on the effect of dietary processed cocoa pod husk in broiler chickens nutrition. Therefore, this study is aimed in determining the effects of fed concentrates supplemented with processed cocoa pod waste on broiler chickens.

## METHODS AND MATERIALS:

### Location of study

The experiment was carried out at the Product Development Unit of the Research Department, University of Uyo, Uyo, Akwa Ibom State. Akwa Ibom state is in Nigeria. It is located in the coastal southern part of the country, lying between latitudes 4°32'N and 5°33'N, and longitudes 7°25'E and 8°25'E. The state is located in the south-South geographical zone, and is bordered on the east by Cross River State and Rivers State, on the west by Abia state, and on the south by Atlantic Ocean and the south-most tip of Cross Rivers State.

### Collection and preparation of test ingredients

Cocoa pod waste were collected from the cocoa farm in Uyo cash crop farm. Later the husk were chopped into slices at the Product Development Unit Of the Research Department. There were sundried for about 24 hours to reduce the moisture content to about 80%. The pre-dried slices were then passed through a combination mincer and pelleting machine to produce pellets. The pellets were again dried for about 4 days to further reduce the moisture content to about 10% and stored until used.

### Experimental diets, Birds, Housing and experimental Design

Three experimental diets were formulated to meet the minimum requirements of the birds at both the starter and finisher phase, in which PCHM was included at 0, 5 and 10% and designated as diets 1, 2 and 3 respectively. One hundred and eighty 1-day-old Arbor Acres broiler chicks were randomly distributed to three dietary treatments (10 birds/replicate; 60 birds/treatment) in a completely randomized design (CRD). The birds in each replicate were housed in their respective wood shavings littered 200 × 100 cm pen. The experimental house temperature was maintained within 31 °C ± 2 for the first 7 days and reduces by 2 °C after each consecutive 7 days until the house temperature was 26 °C ± 2. Illumination was provided for 23 h/day. The birds were fed water and mash ad libitum throughout the experimental period.

### Statistical analysis:

The data were subjected to one-way analysis of variance using SPSS version 20. The differences among means were determined by Duncan multiple range test of the same package.

## RESULT AND DISCUSSION:

Table 1 shows the proximate composition and phytochemicals in the PCHM.

**Table 1: Chemical composition of processed cocoa pod husk**

I n g r e d i e n t s	Q u a n t i t y ( g / k g )
A s h ( % )	1 5 0 . 3 0
C r u d e f i b r e ( % )	1 4 7 . 6 2
C r u d e p r o t e i n ( % )	1 3 7 . 1 3
E t h e r E X t r a c t ( % )	6 4 . 3 8
N i t r o g e n f r e e E x t r a c t ( % )	3 8 2 . 1 2
C a f f e i n e	0 . 0 3
T a n i n	0 . 1 1
T h r e b o m i n e	0 . 3 5
M E	2 4 2 1 . 9 8 K c a l / k g

Metabolizable = (37 × %CP) + (81.8 × %FAT) + (35.5 × %NFE) Pauzenga[17], ME metabolizable

**Table 2: Composition of Experimental Diet**

I n g r e d i e n t s ( % )	Diet 1, 0% PCHM	Diet 2, 5% PCHM	Diet 3, 10% PCHM
M a i z e	4 0 . 0 0	4 0 . 0 0	4 0 . 0 0
S B M	1 0 . 2 3	1 3 . 7 5	2 7 . 5 0
F i s h m e a l	2 5 . 6 0	2 6 . 7 8	2 7 . 9 2
B o n e M e a l	1 . 5 0	1 . 5 0	1 . 5 0
O y s t e r s h e l l	2 . 0 0	2 . 0 0	2 . 0 0
P a l m O i l	1 . 0 0	2 . 0 0	2 . 2 0
P r e m i x *	0 . 2 5	0 . 2 5	0 . 2 5
M e t h i o n i n e	0 . 3 5	0 . 3 5	0 . 3 5
L y s i n e	0 . 2 0	0 . 2 0	0 . 2 0



L e v e l s o f i n c l u s i o n ( % )				
	0 (control)	1	2	1 4
P a r a m e t e r s	1	2	3	S E M ( ± )
Slaughter weight (g/ bird)	2 4 2 4 . 7 1 <sup>a</sup>	2 5 6 2 . 4 9 <sup>a b</sup>	2 8 3 7 . 4 8 <sup>b</sup>	1 2 6 . 3 3
Dressed weight (g/ bird)	1 8 7 1 . 3 2	1 9 3 8 . 0 0	2 1 6 3 . 0 1	9 7 . 4 5
Dressing percentage ( % )	7 4 . 8 7 <sup>a</sup>	7 2 . 4 6 <sup>b</sup>	7 1 . 3 7 <sup>c</sup>	0 . 4 5
I n t e r n a l O r g a n s ( % )				
H e a r t	0 . 3 0 <sup>a</sup>	0 . 3 0 <sup>b</sup>	0 . 3 9 <sup>a</sup>	0 . 0 3
L u n g	1 . 6 3	1 . 5 3	1 . 5 1	0 . 1 2
L i v e r	2 . 0 4 <sup>b</sup>	2 . 2 9 <sup>b</sup>	2 . 6 9 <sup>a</sup>	0 . 1 0
S p l e e n	0 . 2 9 <sup>b</sup>	0 . 4 8 <sup>b</sup>	0 . 4 6 <sup>a</sup>	0 . 1 0
P a n c r e a s	0 . 2 1 <sup>b</sup>	0 . 3 0 <sup>a</sup>	0 . 3 2 <sup>a b</sup>	0 . 2 0
K i d n e y	3 4 . 4 3	3 4 . 8 8	3 4 . 5 2 <sup>a</sup>	0 . 2 0
G i z z a r d	2 . 0 4 <sup>b</sup>	2 . 2 9 <sup>a b</sup>	2 . 6 9 <sup>a</sup>	0 . 2 3
G a l l b l a d d e r	2 . 9 0 <sup>a b</sup>	1 . 8 6 <sup>c</sup>	2 . 0 0 <sup>b c</sup>	0 . 3 7

a,b,c Means in the same row with different super script are significantly different

Nutrition has marked effect on yield of quality meat of animals, and their relative organ weights are very useful in the prediction of toxic effect of the test materials or the diets. In addition, the toxins in diet could be absorbed and accumulated in the various target tissues or organs and cause injury to the cells and alter their normal structure or function. The similarity in the carcass traits and relative internal organ weights of the experimental birds fed the varying inclusion levels of PCHM is of health benefits and indicates that the phytochemicals in PCHM is or fatal effects or that the dietary treatment in this study did not pose treats to the development of edible portion of the experimental birds and the normal gross anatomy of their internal organs.

The carcass and internal organ characteristics of chicken broilers with processed cocoa pod husk meals shown in table 5. Significant differences were found for percentage slaughter weight ( $P<0.05$ ) and also for dressing percentage ( $P>0.05$ ) showing that, the replacement of maize with processed cocoa pod husk meal in the diets of finisher chicken broilers did affect the resulting cold carcass weights of the chicken broilers when slaughtered. This was not in agreement with the findings of Johri [23], Alagbe[24], who also reported that, multi-enzyme addition did not affect carcass characteristics of broilers used in their study.

**Table 6: Effects of processed cocoa pod waste meal on the erthrogram (i.e haemoglobin concentration/ serum biochemistry indices**

P a r a m e t e r s	A	B	C	S E M
H a e m a t o l o g i c a l I n d i c e s				
Packed cell Volume ( % )	3 3 . 0 8	3 1 . 6 3	3 1 . 8 8	2 6 6
Haemoglobin concentration(g/l)	1 0 . 0 2	1 0 . 5 0	1 0 . 7 0	0 0 5
Red Blood Cell ( $\times 10^6$ /UL)	2 . 5 6	2 . 5 9	2 . 6 9	0 2 7
S e r u m b i o c h e m i c a l i n d i c e s				
Total protein ( mg / dl )	1 3 1 . 9 8 <sup>a</sup>	1 2 5 . 1 8 <sup>b</sup>	1 2 6 . 8 5 <sup>a b</sup>	0 2 3
C h o l e s t e r o l ( m m o l / l )	3 8 . 8 8 <sup>a</sup>	3 8 . 8 8 <sup>b</sup>	3 7 . 5 3 <sup>b</sup>	0 0 9
Alanine amino transferase ( % )	2 9 . 0 3 <sup>b</sup>	3 1 . 0 2 <sup>a</sup>	2 9 . 3 3 <sup>b</sup>	2 1 0
Aspartate amino transferase ( U/l )	2 2 . 6 7 <sup>a</sup>	2 0 . 8 3 <sup>a</sup>	1 8 . 3 3 <sup>a</sup>	5 1 5
S e r u m a n t i o x i d a n t e n z y m e				
Glutathione (Mmol/GSSG/min)	7 7 . 6 7 <sup>c</sup>	8 3 . 8 3 <sup>a b c</sup>	8 1 . 0 0 <sup>b c</sup>	0 5 9
C a t a l a s e ( K u / m l )	2 2 . 3 3 <sup>a</sup>	1 6 . 1 7 <sup>a b c</sup>	1 9 . 0 0 <sup>a b</sup>	0 . 0 3 1

a,b,c Means in the same row with different super script are significantly different

Erythrogram is one of the indicators for assessing the nutritional and health status of animals, and there exists a marked influence of nutrition on haematology traits [25]. The stability of packed cell volume, haemoglobin concentration and red blood cells of the birds fed diets containing varying levels of PCHM also shows that the dietary treatment used in this study did not have negative effects on the normal blood-forming processes in the experimental birds. This result agrees with Adeyeye et al. [2], who reported similar haematological indices values among experimental rabbits fed processed cocoa pod husk meal-inclusive diets. The assessment of biochemical parameters is also another important method of assessment of health in animals [26]. The non-difference in the serum biochemical indices values in broiler chicken fed the experiment diets also indicates that dietary PCHM inclusion up to 8% support normal health in the broiler chickens. This may be the product of activities of the phytochemicals in the PCHM. For instance, caffeine intake

was associated with a lower risk of elevated alanine aminotransferase [17]. The use of phytochemical in ameliorating the negative effects of heat-induced oxidative stress in birds has been reported [25].

Table 6 shows the effect of PCHM on haemato-biochemical indices and serum antioxidant enzymes. The packed cell volume, haemoglobin concentration and red blood cells of the broiler chickens were stable ( $P > 0.05$ ) across the dietary treatment. In the same vein, the serum biochemical indices concentration in the experimental birds was not affected ( $P > 0.05$ ) by the dietary treatment. The serum GPx and CAT concentration were higher ( $P < 0.05$ ) in birds broilers fed PCHM-inclusive diets compared to those fed the control diet.

## CONCLUSION:

This study was conducted to determine the growth performance and economic impact of broiler chicken fed concentrates supplemented with processed cocoa pod waste meal. A total of 180 1-day-old Arbor Acres broiler chicks were randomly distributed to three dietary treatments (10 birds/replicate; 60 birds/treatment) in a completely randomized design (CRD) and used for the study. Three experimental diets were formulated to meet the minimum requirements of the birds at both the starter and finisher phase, in which PCHM was included at 0, 5 and 10% and designated as diets 1, 2 and 3 respectively. The experimental house temperature was maintained within  $31^{\circ}\text{C} \pm 2$  for the first 7 days and reduces by  $2^{\circ}\text{C}$  after each consecutive 7 days until the house temperature was  $26^{\circ}\text{C} \pm 2$ . Illumination was provided for 23 h/day. The birds were fed water and mash ad libitum throughout the experimental period the data were subjected to one-way analysis of variance using SPSS version 20 while the differences among means were determined by Duncan multiple range test of the same package. The following were the concluded:

Various levels of inclusion of processed cocoa pod waste(husk) meal was significant with the growth performance, economic evaluation, and growth performance of pigs ( $P = 0.05$ ). Dietary treatment had effect on the feed conversion ratio and feed cost per unit weight gain. Birds fed the control diet (0 PCHM) were lowest in dressing percentage whereas counterparts on 10% maize replacement with processed cocoa pod husk meal (PCHM) where highest in abdominal fat compared to pigs on other dietary treatments. This shows that feeding PCHM aid in internal organs increments and therefore results in significantly increment in carcass fat. Results from haematology and serum biochemical analysis shows that PCHM had no untoward effect on physiology and metabolism to the pigs.

The findings of this study therefore implies that cocoa pod waste meal supplemented can be a great meal in for all monogastric animals as it will increase the body weight, aid growth and promote large meat production. Also, it will reduce the cost expenses on the farmers as the cost of making or getting this feed is relatively easy and low. It is therefore recommended that processes cocoa pod waste meal;

1. Should be encouraged in the feeding of broiler chickens to reduce over dependence of maize feeds by our farmers which have led to high cost of raising monogastrics.
2. Public extension/ advisory staff should be mobilized to convey these results to practicing farmers.
3. The inclusion of PCHM as shown in the result above promote the growth performance and feed conversion efficiency of broiler chicken and as such as been recommended for use by farmers.

## REFERENCES

1. Nworgu, F. C., Adebawale, E. A., Oredein, O. A. & Oni, A. (1999) Prospect and economics of broiler production using two plant protein sources. *Tropical Journal of Animal Science* 2,159–166
2. Adeyeye, S. A., Agbede, J. O., Aletor, V. A., & Oloruntola, O. D. (2017) Processed cocoa (Theobroma cacao) pod husks in rabbits diet: effect on haematological and serum biochemical indices. *Asian Journal of Advanced Agricultural Research* 2(4),1–9
3. Egbunik, G. N., Agiang, E. A., Owosibo, A., & Fatufe, A. A. (2009) Effect of protein on performance and haematology of broiler fed cassava peel-based diets. *Architectural Zootechnology*, 58(224),655–662
4. Ogunsipe, M. H., Ibidapo, I., Oloruntola, O. D. & Agbede, J. O. (2017a) Growth performance of pigs on dietary cocoa bean shell meal. *Livestock Res Rural Dev* 29(1) .
5. Ogunsipe, M. H., Balogun, K. B., Oladepo, A. D., Ayoola, M. A. & Arikewuyo, M. T. (2017b) Nutritive value of cocoa bean shell meal and its effect on growth and haematology of weaning rabbits. *Nigerian Journal of Agricultural Food Environment* 13(1),23–28
6. Oloruntola, O. D., Agbede, J. O., Onibi, G. E., Igbasen, F. A., Ayodele, S. O., Arogunjo, S. O. & Ogunjo, S. T. (2018a) Rabbits fed fermented cassava starch residue I: effect on performance and health status. *Architectural Zootechnology*67(260),578–586
7. Oloruntola OD, Agbede JO, Onibi GE, Igbasan FA, Ogunsipe MH, Ayodele SO (2018b) Rabbits fed fermented cassava starch residue II: enzyme supplementation influence on performance and health status. *Architectural Zootechnology*67(260),588–595



8. Akbarian, A., Michiels, J., Degroote, J., Majdeddin, M., Golian, A. & Dmegt, S. D. (2016) Association between heat stress and oxidative stress in poultry: mitochondrial dysfunction and dietary interventions with phytochemicals. *Journal of Animal Science Biotechnology* 7,37.
9. Eghosa, O. U, Rasheed, A. H., Martha, O. & Luqman, A. A. (2010) Utilization of cocoa pod husk (CPH) as substitute for maize in layers mash and perception of poultry farmers in Nigeria. *International Journal of Science Nature* 1(2),271–275
10. Gemed, H. F., & Ratta, N. (2014) Antinutritional factors in plant foods: potential health benefits and adverse effects. *International Journal of Nutritional Food Science*, 3(4),284–289  
<http://www.sciencepublishinggroup.com/ijnf>  
<https://doi.org/10.1186/s40104-016-0097-5>
11. Alemawor, F., Dzogbefia, V. P., Oddoye, E. K. & Oldham, J. H. (2009) Effect of *Pleurotus ostreatus* fermentation on cocoa pod husk composition: influence offermentation period and Mn<sup>2+</sup> supplementation on the fermentationprocess. *African Journal of Biotechnology* 8(9),1950–1958
12. Oloruntola, O. D., Agbede, J. O., Onibi, G. E., Igbasan, F. A. (2015) Composition of cassava (*Manihot* spp.) peels fermented with bovine rumen liquor and different nitrogen sources. *Journal of Global Agricultural Ecology* 2(1),26–35
13. Adamafo, N. A., Cooper Aggrinage. E., Onaye, E. O., Laary, J. K., & Onaye, J. (2004) Effectiveness of corn stalk ash in reducing tannin level and improving in vitro enzymatic degradation of polysaccharides in crop residues. *Ghana Journal of Science*, 44,87–92
14. Oloruntola, O.D., Ayodele, S. O. & Oloruntola, D. A. (2018c) Effect of pawpaw (*Carica papaya*) leaf meal and dietary enzymes on broiler performance, digestibility,carcass and blood composition. *Review Elevated Medical Veterinary Pays Tropics* 71(3). <https://doi.org/10.19182/remvt.31640>
15. Adeyeye, S. A., Agbede, J. O., Aletor, V. A. & Oloruntola, O. D. (2018) Performance and carcass characteristics of growing rabbits fed diets containing graded levels of processed cocoa (*Theobroma cacao*) pod husk meal supplemented with multi-enzyme. *Journal of Applied Life Science International*, 17(2),1–11
16. Pazenga U (1985) Feeding Parent Stock. Zoo Technical International. pp 22–24.
17. Ruhl, C. E, & Everhart, J. E. (2005) Coffee and tea consumption are associated with a lower incidence of chronic liver disease in the United States. *Gastroenterology* 129,1928–1936
18. Parvin N, Mandal TK, Saxema V, Sarkar S, Saxena AK (2010) Effect of increasing protein percentage feed on the performance and carcass characteristics of broiler chicks. *Asian Journal of Poultry Science* 4(2),53–59
19. Akinfala, E. O., Aderibigbe, A. O., & Matanmi, O. (2002) Evaluation of the nutritive value of whole cassava plant as replacement for maize in the starter diets for broiler chickens. *Livestock Research for Rural Development* 14(6). <http://www.lrrd.org/lrrd14/6/akin146.htm>
20. Afolayan, S. B., Dafwang, I. I., Tegbe, T. B. & Sekoni, A. (2012) Response of broiler chickens fed maize-based diets substituted with graded levels of sweet potato meal. *Asian Journal of Poultry Science* 6(1),15–22
21. Gajana, C. S., Nkukwana, T. T., Chimonyo, M., & Muchenje, V. (2011) Effect of altering the starter and finisher dietary phases on growth performance of broilers. *African Journal of Biotechnolgy* 10(64),14203–14208  
<http://www.academicjournals.org/AJB>
22. Ferket, P. R., & Gernat, A. G. (2006) Factors that affect feed intake of meat birds: a review. *International Journal of Poultry Science*, 5(10),905–911
23. Johri, M. O. (2004), Effect of pawpaw (*Carica papaya*) leaf meal and dietary enzymes on broiler performance, digestibility,carcass and blood composition. *Review Elevated Medical Veterinary Pays Tropics* 71(3).
24. Alagbe S. T.(2017) Effect of altering the starter and finisher dietary phases on growth performance of broilers. *African Journal of Biotechnolgy* 10(64),14203–14208
25. Oloruntola, O. D., Agbede, J. O., Ayodele, S.O. & Oloruntola, D.A. (2018d) Neem, pawpaw, and bamboo leaf meal dietary supplementation in broiler chickens: effect on performance and health status. *Journal Food of Biochemistry*, 12, 723 <https://doi.org/10.1111/jfbc.12723>
26. Milner, J. M., Stien, A, Justin Irvine, R., Albon, S. D., Langvatin, R.& Ropstad, E. (2003) Body condition in Svalbard reindeer and the use of blood parameters as indicator of condition and fitness. *Canada Journal of Zoology* 81,1566–1578